

ATOMIC AND MOLECULAR IMPURITIES IN ^4He CLUSTERS

F. Dalfovo

Dipartimento di Fisica, Università di Trento, 38050 Povo, Italy

Abstract. *A density functional theory is used to predict the binding energy of atomic and molecular impurities (Ne, Ar, Kr, Xe, Li, Na, K, Rb, Cs, and SF_6) in the center of ^4He clusters, in the limit of zero temperature and for zero angular momentum states. The size dependence of the binding energy, from small clusters to the bulk liquid limit, is investigated. The behaviour of the ^4He density near the impurity is also studied.*

PACS: 36.40; 67.40

I. INTRODUCTION

Helium clusters are produced in supersonic nozzle beam expansions (see Ref.[1] for a recent review). They are expected to behave like liquid droplets down to zero temperature. At low temperature they are strongly affected by quantum correlations and, eventually, become superfluid. Several attempts have been made in the last years in order to get experimental evidence of superfluidity in helium clusters. Due to the weakness of the helium-helium interaction a direct characterization of pure helium clusters is very difficult. A more promising approach is the investigation of atomic and molecular impurities attached to the helium clusters [2,3]. The interpretation of the experimental data is still limited by the lack of quantitative theories. The quantum mechanical description of the static and dynamics of helium clusters has been tackled by several authors [4-10], but very little is known about the behaviour of impurities [9-11].

In this work we calculate the energy and the density distribution associated with impurity states in the center of helium clusters at zero temperature and zero angular momentum. We employ a density functional method, which was developed in the last decade [5,12-14] in the context of inhomogeneous states of liquid helium. With a relatively small numerical effort it provides quantitative predictions which are close to the results of *ab initio* Monte Carlo calculations in the case of small clusters, and can be extended to large clusters, up to the bulk liquid limit. Quantum correlations between helium atoms are accounted for by means of a phenomenological density dependent interaction. The impurity is included as an external potential in which the helium density adjusts to minimize the energy. Rare gas atoms, alkali atoms, and SF₆ molecule are considered. Accurate impurity-helium potentials are taken from the literature [15-18].

In Section II we introduce the density functional formalism, emphasizing the main physical features and discussing the approximations made in the treatment of the impurity. In Section III we present the results of the calculations. Section IV is a short summary of the main results and a discussion about open problems and future work.

II. METHOD

In a density functional theory one writes the total energy of the many-body system as a functional of the one-body density $\rho(\mathbf{r})$, in the form

$$E = \int d\mathbf{r} \mathcal{H}[\rho] \quad . \quad (1)$$

Under certain conditions the minimization of E with respect to ρ is equivalent to the solution of the many-body Schrödinger equation [19]. In general, however, the exact form of the functional, whose minimum is located at the true equilibrium density of the system, is not known *a priori*. Thus suitable phenomenological functionals are introduced; they yield approximate results, whose quality depends on how the relevant symmetries and correlations are included in the starting functional form.

A first systematic description of helium clusters in the framework of density functional is the one of Ref. [5]. The theory was built in such a way that the compressibility and surface tension of liquid helium were reproduced. The results for the cluster energy and density profile were close to the ones of quantum variational calculations [4]. The functional of Ref. [5] has been recently extended to include the effect of the finite range helium-helium interaction [13]. The new functional has the form:

$$\mathcal{H}_o = \frac{\hbar^2}{2m} (\nabla \sqrt{\rho(r)})^2 + \frac{1}{2} \int d\mathbf{r}' \rho(r) \rho(r') V(|\mathbf{r} - \mathbf{r}'|) + \frac{c}{2} \rho(\mathbf{r}) (\bar{\rho}_{\mathbf{r}})^{1+\gamma} \quad . \quad (2)$$

The first term in the sum is a quantum pressure; it corresponds to the zero temperature kinetic energy of a noninteracting Bose system. The second term contains a two-body interaction V , which is the Lennard-Jones interatomic potential, with the standard parameters $\alpha = 2.556 \text{ \AA}$ and $\varepsilon = 10.22 \text{ K}$, screened at a distance $h = 2.377 \text{ \AA}$ with a power law, as follows

$$V(x) = \begin{cases} 4\varepsilon \left[\left(\frac{\alpha}{x} \right)^{12} - \left(\frac{\alpha}{x} \right)^6 \right], & \text{if } x \geq h; \\ V(h) \left(\frac{x}{h} \right)^4, & \text{if } x < h. \end{cases} \quad (3)$$

The last term in Eq. (2) accounts for short range correlations between atoms. In particular it contains the effect of the hard core part of the interatomic potential. Its form follows the idea of the "weighted density approximation", used mainly in the study of classical fluids. The weighted density $\bar{\rho}$ is the average of $\rho(\mathbf{r})$ over a sphere with radius h . The parameters $c = 1.04554 \times 10^7 \text{ K \AA}^{3(1+\gamma)}$, and $\gamma = 2.8$, together with the screening length h , are the only three parameters of the theory; they are fixed to reproduce the equation of state of the bulk liquid. As discussed in ref.[13] the functional $\mathcal{H}_o$ corresponds to a mean-field description, which incorporates phenomenologically the effects of a finite-range interaction, with the correct long range behaviour, as well as of short range correlations. Like the functional of Ref. [5] it reproduces the equation of state, the compressibility and the surface tension of bulk liquid ^4He . Moreover it reproduces the behaviour of the static response function, which is peaked at the roton wavelength. This ensures the inclusion of localization effects, which are crucial to predict the freezing transition at high pressure [20], the layer structure in helium films [21,22], and the shell structure near impurities [14].

The effect of an impurity can be included in the density functional by adding a potential term

$$\mathcal{H} = \mathcal{H}_o + V_I(\mathbf{r})\rho(\mathbf{r}) \quad , \quad (4)$$

where $V_I(\mathbf{r})$ is the helium-impurity potential. This corresponds to treat the impurity

as a classical object in a quantum liquid. Since the zero point motion of the impurity is not accounted for, the validity of Eq. (4) is restricted to impurities heavier than the helium atomic mass m . So, we do not discuss ^3He or hydrogen impurities. For them a full quantum mechanical treatment is necessary [10,11]. In this work we consider rare gas atoms, alkali atoms and SF_6 molecules. For most of them the effect of zero point motion is surely negligible. Only for Li and Ne it gives rise to sizeable effects, but, as we will discuss later, the potential contribution is expected to be still dominant.

Combining Eqs. (1), (2) and (4) the minimization with respect to ρ yields the following Euler-Lagrange equation:

$$\left[-\frac{\hbar^2}{2m}\Delta + U(\mathbf{r}) + V_I(\mathbf{r}) \right] \sqrt{\rho} = \mu\sqrt{\rho} \quad , \quad (5)$$

where μ is the helium chemical potential, while $U(\mathbf{r})$ is the Hartree self consistent field derived from functional (2) [13,14]. Equation (5) can be solved in bulk helium as well as in clusters of given particle number N . The solution yields both the density profile and the energy of the system. The impurity chemical potential can be extracted as difference between the energy of the cluster with and without the impurity (X)

$$\mu_I = E[\text{X}(\text{He})_N] - E[(\text{He})_N] \quad . \quad (6)$$

In bulk this is equivalent to

$$\mu_I = \int d\mathbf{r} \quad (\mathcal{H}[\rho(\mathbf{r})] - \mu\rho(\mathbf{r})) \quad . \quad (7)$$

The solution of Eq. (5) in bulk helium as been already discussed by Pavloff [14], who considered Na, Cs, Xe and Ba^+ as impurities and used the same density functional for liquid helium. The results for Xe were in good agreement with previous variational calculations by Kürten and Ristig [23]. For Cs the agreement was

only qualitative, mainly because the role of the elementary diagrams neglected in the calculations of Ref. [23] is expected to be more important for Cs than for Xe. Another test on the density functional method is the evaluation of the energy of one atom of ^4He , considered as a classical impurity in the rest of the liquid. The calculation, with helium-helium interaction in place of V_I , yields an energy of about -23 K. This is in good agreement with independent estimates of the potential energy per particle in bulk helium. The difference between this value and the total energy per particle, -7.15 K, is an estimate of the zero point kinetic energy. The kinetic energy for different impurities scales approximately as the mass ratio and the square of the radius of the bubble created by the impurity inside the liquid. For the lightest impurities considered in this work (Li and Ne) the resulting zero point energy is no more than $3 \div 4$ K, which is much less than the potential energy.

In the present work we calculate the impurity chemical potential and the helium density near the impurity, in bulk liquid and clusters, using accurate impurity-helium interaction [15-18]. We consider only impurity states in spherical symmetry. Thus Eq. (5) becomes a one-dimensional equation in the radial coordinate, which can be solved numerically with a standard iterative procedure (a few minutes on a RISC-CPU for each run). In the case of helium clusters, the assumption of spherical symmetry implies impurity states in the center of the cluster. The inclusion of possible bound states outside the cluster (surface states) will be a subsequent step, which requires further numerical efforts in solving Eq. (5) with anisotropic density.

III. RESULTS

III.A Impurity in Bulk Liquid

We take the mixed rare gas van der Waals potential by Tang and Toennies [16], the alkali-helium potential by Patil [18], and the spherical approximation of the SF₆-helium potential by Pack et al. [15], with a modified value of the potential depth as in Ref. [17]. We solve Eq. (5) in bulk liquid helium by imposing the asymptotic value of ρ equal to the uniform liquid density at fixed external pressure. The results for the impurity states are summarized in Table I. In the first column the impurity chemical potential (at zero temperature and zero pressure) is given. One notes that it is negative for rare gas atoms and for SF₆, while it is positive for alkali atoms. This reflects the different structure of V_I ; the alkali-helium potential has a more extended repulsive core and a weaker attractive tail than the rare-gas and the SF₆ potential. As a consequence the rare gas impurities, as well as SF₆, tend to compress the helium atoms in shells around the impurity, in a region of large and negative V_I , with a gain in energy. On the contrary, the alkali atom pushes the helium atoms far away without changing significantly the helium density; this corresponds to a cost in energy. Density profiles are shown in Fig. 1. The height of the first peak in helium density is also given in the second column of Table I (the equilibrium density of the uniform liquid is $0.0218 \text{ \AA}^{-3}$). The energy systematics and the structure of the density profiles are very similar to the situation of helium films on solid substrates [21,22], where one finds a transition from wetting to non wetting as a function of the substrate-helium potential parameters.

An interesting quantity is the number of helium atoms in the first shell, close to the impurity (third column in Table I). The most attractive case is the one of SF₆, where we find approximately 25 atoms in the first shell, corresponding to a

solid *snowball* surrounding the impurity. This *snowball* makes the impurity state less sensitive to the external pressure, since the behaviour of state is dominated by the local shell structure. The chemical potential of alkali atoms is much more pressure dependent. It tends to increase with the external pressure, because it costs more energy to create a *bubble* in a liquid under pressure. For instance, the chemical potential of Na increases from 49 K to 74 K in passing from zero to 5 bars, while the one of Kr decreases from -243 K to -250 K in the same range. It is worth noticing that the existence of a shell structure near an impurity, as in the case of SF_6 and rare gas atoms, has important consequences in the dynamic properties of the system. In particular it is expected to change considerably the effective mass of the impurity as well as the structure of the velocity field of helium in the surrounding region.

The accuracy of our predictions depends on two main aspects: the quality of the impurity-helium interaction used as input and the quality of the density functional $\mathcal{H}_0$. Indeed one can take from the literature different impurity-helium potential for the same kind of impurity. The resulting chemical potential scales approximately as the well depth of V_I , and the height of the first peak in ρ slightly changes. But, at the present level of understanding, the precision in the impurity-helium potential is not crucial. For this reason we have also chosen the isotropic approximation for the SF_6 -helium potential instead of the true anisotropic one. As concerns the quality of the density functional, one can test it by comparison with quantum Monte Carlo calculations, whenever available, as done in different contexts [5,11,13,22,24].

III.B Impurities in the center of helium clusters

The solution of Eq. (5) without the impurity potential corresponds to the case

of pure ${}^4\text{He}$ clusters. The results for the energy systematics and for the density profiles are very close to the ones of Ref. [5], where a simpler density functional was used. This confirms the fact that for smoothly varying density the relevant quantities in the theory are the compressibility and the surface tension, which are both well reproduced by the two functionals.

We solve Eq. (5) with and without impurities. The difference in energy provides the impurity chemical potential. A major advantage of the density functional method is that it works easily even with quite large clusters, up to $N = 5000$ particles or more. So one can test the asymptotic convergence to the bulk liquid limit, predicting the size dependence of the results on a wide range of N .

In Figs. 2 and 3 we show the impurity chemical potential as a function of N . Again there is a significant difference between alkali atoms and the other impurities. The alkali atoms are more sensitive to the size of the cluster and reach the asymptotic value of the chemical potential in bulk more slowly. This effect can be understood by looking at Fig. 4, where the density profile for a cluster with Na and Kr is compared with the pure helium cluster. One notes that the rare gas atom does not modify the external structure of the cluster. As a consequence its chemical potential is fixed mainly by the local distortion of the density near the center, and the energy is almost the same as in bulk liquid. On the contrary the Na atom pushes the helium atoms outside, working against the local pressure of the liquid and increasing the surface area. The net effect is an increase of the impurity chemical potential with respect to the bulk value. This is also in agreement with the pressure dependence of the chemical potential in bulk, which is stronger for alkali atoms.

In all cases the chemical potential bends towards zero in the limit of small clusters. This happens when the first shells of atoms near the impurity become

partially occupied. In the case of SF_6 and rare gas the attractive impurity-helium potential dominates on the helium-helium correlations; thus a lack of atoms in the first shells, in the potential well of the impurity, implies an increase of energy. Viceversa in the case of alkali atoms the helium-helium correlations are more important and a decrease in helium density implies a decrease in energy.

Typical density profiles are shown in Fig. 5. One notes that the first shell of atoms near the rare gas impurity is deformed only for very small N . This is a clear sign that the impurity prefers to tie the helium atoms around itself, i.e., to stay in the center of the cluster with a relatively large binding energy. This effect is even stronger in the case of SF_6 . On the contrary, the alkali atoms seem not to bind to the center of the cluster for any value of N . The energy of the impurity-cluster system is higher when the impurity is in the center than when it is far outside. This does not exclude *a priori* the possibility to capture alkali atoms on helium droplets. In fact, since at large distance the relative interaction is attractive, a local energy minimum may still exist on the surface of the cluster.

Monte Carlo calculations for small clusters with impurities are also becoming available. In Fig. 6 we compare the density profile for a cluster of 111 particles with a SF_6 molecule in the center. The dashed line is the result of Diffusion Monte Carlo calculations [10], and the solid line is the prediction of our density functional theory, using the same impurity-helium potential. The agreement is very good. The small difference in the first peak height is well within the expected accuracy of the present theory.

IV. CONCLUSIONS

We have done density functional calculations for rare gas atoms, alkali atoms

and SF_6 molecule in the center of helium clusters. We have discussed the size dependence of the impurity chemical potential and of the helium density profile on a wide range of particle number, up to the bulk liquid limit. Our results strongly support the existence of bound states in the center of a cluster for rare gas atoms and SF_6 , but not for alkali atoms. For SF_6 in small clusters the predictions of the density functional theory are in good agreement with the ones of recent Diffusion Monte Carlo calculations [10].

In view of the current debate about the location of impurities on clusters [2,3] further theoretical work is needed. A first possibility is to drop the spherical symmetry in Eq. (5), to allow for impurity states on the surface of the clusters, as well as to study the cluster-impurity interaction as a function of the relative distance. This makes the numerical computation heavier, without any substantial change in the theory. A second point is the inclusion of non zero angular momentum, which is expected to favour surface impurity states [2]. The inclusion of velocity field and vorticity in the framework of the density functional theory has been already discussed for bulk liquid helium [24]. The treatment of the velocity field in the cluster geometry is expected to be more difficult. The problem deserves certainly further investigations, being directly related to the concept of superfluidity in finite systems.

ACKNOWLEDGEMENTS

This work was partially supported by INFN, gruppo collegato di Trento.

REFERENCES.

1. Toennies, J.P.: In: *Proceedings of the International School of Physics E. Fermi on "Chemical Physics of Atomic and Molecular Clusters", Course CVII*, p.597. Scoles, G. (ed.). Amsterdam: North Holland 1990
2. Schutt, D.L.: Thesis, Princeton 1992; Goyal, S., Schutt, D.L., Scoles, G.: Phys. Rev. Lett. **69** 933 (1992); Goyal, S., Schutt, D.L., Scoles, G.: Chem. Phys. Lett. **196**, 123 (1992); Goyal, S., Schutt, D.L., Scoles, G.: J. Phys. Chem. **97**, 2236 (1993)
3. Scheidemann, A., Toennies, J.P., Northby, J.A.: Phys. Rev. Lett. **64**, 1899 (1990); Scheidemann, A., Schilling, B., Toennies, J.P.: Preprint, Göttingen 1993
4. Pandharipande, V.R., Zabolitzky, J.G., Pieper, S.C., Wiringa, R.B., Helbrecht, U.: Phys. Rev. Lett. **50**, 1676 (1983); Pandharipande, V.R., Pieper, S.C., Wiringa, R.B.: Phys. Rev. B **34**, 4571 (1983); Pieper, S.C., Wiringa, R.B., Pandharipande, V.R.: Phys. Rev. B **32**, 3341 (1985)
5. Stringari, S., Treiner, J.: J. Chem. Phys. **87**, 5021 (1987)
6. Casas, M., Stringari, S.: J. Low Temp. Phys. **79**, 135 (1990)
6. Sindzingre, P., Klein, M.L., Ceperley, D.M.: Phys. Rev. Lett. **63**, 1601 (1989)
7. Rama Krishna, M.V., Whaley, K.B.: Phys. Rev. Lett. **64**, 1126 (1990); Rama Krishna, M.V., Whaley, K.B.: J. Chem. Phys. **93**, 746 (1990); Rama Krishna, M.V., Whaley, K.B.: J. Chem. Phys. **93**, 6738 (1990)
8. Chin, S.A., Krotscheck, E.: Phys. Rev. Lett. **65**, 2658 (1990); Chin, S.A., Krotscheck, E.: Phys. Rev. B **45**, 852 (1992)
9. Barnett, R.B., Whaley, K.B.: J. Chem. Phys. **96**, 2953 (1992)
10. Whaley, K.B.: Preprint, Berkeley 1993; Barnett, R.B., Whaley, K.B.: (in

preparation)

11. Dalfovo, F.: Z. Phys. **D 14**, 263 (1989)
12. Stringari, S., and Treiner, J.: Phys. Rev. **B 36**, 8369 (1987)
13. Dupont-Roc, J., Himbert, M., Pavloff, N., Treiner, J.: J. Low Temp. Phys. **81**, 31 (1990)
14. Pavloff, N.: Thesis, Orsay 1990
15. Pack, R.T., Piper, E., Pfeffer, G.A., Toennies, J.P.: J. Chem. Phys. **80**, 4940 (1984)
16. Tang, K.T., Toennies, J.P.: Z. Phys. D **1**, 91 (1986)
17. Scoles, G.: Int. J. of Quantum Chemistry: Quantum Chemistry Symposium **24**, 475 (1990)
18. Patil, S.H.: J. Chem. Phys. **94**, 8089 (1991)
19. Hohenberg, P., Kohn, W.: Phys. Rev. **126**, B864 (1964); Kohn, W., Sham, J.: *ibid* **140**, A1133 (1965); Mermin, N.D.: *ibid* **137**, A1441 (1965)
20. Dalfovo, F., Dupont-Roc, J., Pavloff, N., Stringari, S., Treiner, J.: Europhys. Lett. **16**, 205 (1991)
21. Pavloff, N., Treiner, J.: J. Low Temp. Phys. **83**, 331 (1991)
22. Cheng, E., Cole, M.W., Saam, W.F., Treiner, J.: Phys. Rev. Lett. **67**, 1007 (1991); Cheng, E., Cole, M.W., Saam, W.F., Treiner, J.: Phys. Rev. B **46**, 13967 (1992)
23. Kürten, K.E., Ristig, M.L.: Phys. Rev. B **27**, 5479 (1983); Kürten, K.E., Ristig, M.L.: Phys. Rev. B **31**, 1346 (1985)
24. Dalfovo, F.: Phys. Rev. B **46**, 5482 (1992)

FIGURE CAPTIONS

- Fig. 1** Helium density profiles near impurities in bulk liquid, at zero temperature and zero pressure. The profiles for rare gas impurities correspond to Ne, Ar, Kr, and Xe in this order starting from the profile closest to $r = 0$. The profiles for alkali impurities corresponds to Li, Na, K, Rb, and Cs in the same order. The coordinate r is the distance from the impurity.
- Fig. 2** Chemical potential of rare gas atoms, and SF_6 molecule, in helium clusters as a function of the helium particle number.
- Fig. 3** Chemical potential of alkali atoms in helium clusters as a function of the helium particle number.
- Fig. 4** Density profile for 300 helium atoms around Na and Kr impurity. The dashed line is the density profile of a pure helium cluster with the same number of particles.
- Fig. 5** Density profile for helium clusters with Na and Kr impurities. The number of helium particles is 20, 30, 50, 100, 300, 1000, and 3000 starting from the smallest one.
- Fig. 6** Density profile for a cluster of 111 helium atoms and one SF_6 molecule in the center. Dashed line: Diffusion Monte Carlo calculations of Ref. [10]; solid line: present calculation with the same impurity-helium potential.

TABLE CAPTION

Table I. Results for the impurity state in bulk liquid helium at zero pressure and zero temperature. From left to right: impurity type, impurity chemical potential, maximum value of the helium density, and number of helium atoms in the first shell.

TABLE I

	μ_I [K]	$\rho_{\max}$ ($\text{\AA}^{-3}$)	N_{shell}
Li	+ 40	0.0242	—
Na	+ 49	0.0237	—
K	+ 68	0.0227	—
Rb	+ 69	0.0226	—
Cs	+ 84	0.0220	—
Ne	— 39	0.0477	12
Ar	— 195	0.0593	17
Kr	— 243	0.0605	19
Xe	— 313	0.0595	21
SF ₆	— 601	0.0820	25